

On the neutron decay

Francesco Giacosa

UJK Kielce – Goethe U Frankfurt

In collaboration with: Giuseppe Pagliara, INFN& University of Ferrara

Talk based on 1906.10024 and prepared for:

3rd Jagiellonian Symposium on
Fundamental and Applied Subatomic Physics,
UJ Krakow

23-28/6/2019

Outline



1. General discussion of the decay law: non-exp. decays, QZE and IZE
2. Experimental proofs
3. The neutron decay anomaly
4. The Inverse-Zeno-effect as an explanation of the anomaly
5. Conclusions

Part 1: General discussion

Exponential decay law

- N_0 : Number of unstable particles at the time $t = 0$.

$$N(t) = N_0 e^{-\Gamma t}, \quad \tau = 1/\Gamma \text{ mean lifetime}$$

Confirmed in countless cases!

- For a single unstable particle:

$$p(t) = e^{-\Gamma t}$$

is the survival probability for a single unstable particle created at $t=0$.
(Intrinsic probability, see Schrödinger's cat).

For small times: $p(t) = 1 - \Gamma t + \dots$

Basic definitions



Let $|S\rangle$ be an unstable state prepared at $t = 0$.

Survival probability amplitude at $t > 0$:

$$a(t) = \langle S | e^{-iHt} | S \rangle$$

Survival probability: $p(t) = |a(t)|^2$

Rep. Prog. Phys., Vol. 41, 1978. Printed in Great Britain

Decay theory of unstable quantum systems

L FONDA, G C GHIRARDI and A RIMINI

Deviations from the exp. law at short times

Taylor expansion of the amplitude:

$$a(t) = \langle S | e^{-iHt} | S \rangle = 1 - it \langle S | H | S \rangle - \frac{t^2}{2} \langle S | H^2 | S \rangle + \dots$$

$$a^*(t) = \langle S | e^{iHt} | S \rangle = 1 + it \langle S | H | S \rangle - \frac{t^2}{2} \langle S | H^2 | S \rangle + \dots$$


It follows:

$$p(t) = |a(t)|^2 = a^*(t)a(t) = 1 - t^2 \left(\langle S | H^2 | S \rangle - \langle S | H | S \rangle^2 \right) + \dots = 1 - \frac{t^2}{\tau_Z^2} + \dots$$

$$\text{where } \tau_Z = \frac{1}{\sqrt{\langle S | H^2 | S \rangle - \langle S | H | S \rangle^2}} .$$

Note: the quadratic behavior holds for any quantum transition, not only for decays. It is an absolutely general property.

$p(t)$ decreases quadratically (not linearly);
no exp. decay for short times.
 τ_Z is the 'Zeno time'.



Time evolution and energy distribution (1)

The unstable state $|S\rangle$ is not an eigenstate of the Hamiltonian H .
Let $d_S(E)$ be the energy distribution of the unstable state $|S\rangle$.

Normalization holds: $\int_{-\infty}^{+\infty} d_S(E) dE = 1$

$$a(t) = \int_{-\infty}^{+\infty} d_S(E) e^{-iEt} dE$$

In stable limit: $d_S(E) = \delta(E - M) \rightarrow a(t) = e^{-iMt} \rightarrow p(t) = 1$

Time evolution and energy distribution (2)



Breit-Wigner distribution:

$$d_s(E) = \frac{\Gamma}{2\pi} \frac{1}{(E - M)^2 + \Gamma^2 / 4} \rightarrow a(t) = e^{-iM_0 t - \Gamma t / 2} \rightarrow p(t) = e^{-\Gamma t}.$$


The Breit-Wigner energy distribution cannot be exact.

Two physical conditions for a realistic $d_s(E)$ are:

1) Minimal energy: $d_s(E) = 0$ for $E < E_{\min}$

2) Mean energy finite: $\langle E \rangle = \int_{-\infty}^{+\infty} d_s(E) E dE = \int_{E_{\min}}^{+\infty} d_s(E) E dE < \infty$

A very simple numerical example




$$M_0 = 2; E_{\min} = 0.75; \Gamma = 0.4; \Lambda = 3$$

$$d_s(E) = N_0 \frac{\Gamma}{2\pi} \frac{e^{-(E^2 - E_0^2)/\Lambda^2} \theta(E - E_{\min})}{(E - M_0)^2 + \Gamma^2/4}$$

$$d_{BW}(E) = \frac{\Gamma_{BW}}{2\pi} \frac{1}{(E - M_0)^2 + \Gamma_{BW}^2/4}$$

$$\Gamma_{BW}, \text{ such that } d_{BW}(M_0) = d_s(M_0)$$



$$a(t) = \int_{-\infty}^{+\infty} d_s(E) e^{-iEt} dE; \quad p(t) = |a(t)|^2$$

$$p_{BW}(t) = e^{-\Gamma_{BW} t}$$

The quantum Zeno effect

We perform N inst. measurements:


the first one at time $t = t_0$, the second at time $t = 2t_0$, ..., the N -th at time $T = Nt_0$.

$$P_{\text{after } N \text{ measurements}} = p(t_0)^N \approx \left(1 - \frac{t_0^2}{\tau_Z^2}\right)^N = \left(1 - \frac{T^2}{N^2 \tau_Z^2}\right)^N$$

under the assumption that t_0 is small enough.

$$\text{If } N \gg 1 \text{ (at fixed } T\text{): } P_{\text{after } N \text{ measurements}} \approx e^{-\frac{T^2}{N\tau_Z^2}} \approx 1.$$

For large but finite N :
→ slowing down of the decay.




General description of the Zeno and anti-Zeno effects

$$p(t) = e^{-\gamma(t)t} \Rightarrow \gamma(t) = -\frac{1}{t} \ln p(t)$$

Survival probability after a single measurement at time T

$$p(T) = e^{-\gamma(T)T}$$



Survival probability after N measurements:

$$p(\tau)^N = e^{-\gamma(\tau)\tau N} = e^{-\gamma(\tau)T} > e^{-\gamma(T)T} \quad \text{wenn } \gamma(\tau) < \gamma(T) \quad \text{Zeno effect}$$

For $\tau \rightarrow 0$, $\gamma(\tau \rightarrow 0) \rightarrow 0$, $p(\tau)^N \rightarrow 1$

How it is also possible that: $\gamma(\tau) > \gamma(T) \Rightarrow p(\tau)^N = e^{-\gamma(\tau)\tau N} = e^{-\gamma(\tau)T} < e^{-\gamma(T)T}$ **Anti-Zeno-Effekt**

See: P. Facchi, H. Nakazato, and S. Pascazio, Phys. Rev. Lett. **86**, 2699 (2001).



Does the Lifetime of an Unstable System Depend on the Measuring Apparatus? (*)

A. DEGASPERIS (**) and L. FONDA

International Centre for Theoretical Physics - Trieste

G. C. GHIRARDI

The Zeno's paradox in quantum theory

B. Misra and E. C. G. Sudarshan*

Center for Particle Theory, University of Texas at Austin, Austin, Texas 78712

(Received 24 February 1976)

letters to nature

.....
Acceleration of quantum decay processes by frequent observations

A. G. Kofman & G. Kurizki

NATURE | VOL 405 | 1 JUNE 2000 |

Francesco Giacosa

QUANTUM ZENO EFFECTS WITH "PULSED" AND "CONTINUOUS" MEASUREMENTS

P. Facchi⁽¹⁾ and S. Pascazio⁽²⁾

INTERNATIONAL CONFERENCE

TAQMSB

TIME'S ARROWS, QUANTUM MEASUREMENT AND SUPERLUMINAL BEHAVIOR

Part 2: Experimental evidence of non-exponential decay

Experimental confirmation of non-exponential decays (1)

NATURE | VOL 387 | 5 JUNE 1997


Experimental evidence for non-exponential decay in quantum tunnelling

Steven R. Wilkinson, Cyrus F. Bharucha, Martin C. Fischer, Kirk W. Madison, Patrick R. Morrow, Qian Niu, Bala Sundaram* & Mark G. Raizen


Department of Physics, The University of Texas at Austin, Austin, Texas 78712-1081, USA

An exponential decay law is the universal hallmark of unstable systems and is observed in all fields of science. This law is not, however, fully consistent with quantum mechanics and deviations from exponential decay have been predicted for short as well as long times¹⁻⁸. Such deviations have not hitherto been observed experimentally. Here we present experimental evidence for short-time deviation from exponential decay in a quantum tunnelling experiment. Our system consists of ultra-cold sodium atoms that are trapped in an accelerating periodic optical potential created by a standing wave of light. Atoms can escape the wells by quantum tunnelling, and the number that remain can be measured as a function of interaction time for a fixed value of the well depth and acceleration. We observe that for short times the survival probability is initially constant before developing the characteristics of exponential decay. The conceptual simplicity of the experiment enables a detailed comparison with theoretical predictions.

Cold Na atoms in a optical potential



$$V(x,t) = V_0 \cos(2k_L x - k_L a t^2)$$




$$x' = x - \frac{1}{2} a t^2$$

$$U(x') = V_0 \cos(2k_L x') + Max'$$

Experimental confirmation of non-exponential decays (2)

Measured survival probability $p(t)$



Experimental confirmation of non-exponential decays and Zeno /Anti-Zeno effects



VOLUME 87, NUMBER 4

PHYSICAL REVIEW LETTERS

23 JULY 2001

Observation of the Quantum Zeno and Anti-Zeno Effects in an Unstable System

M. C. Fischer, B. Gutiérrez-Medina, and M. G. Raizen

Department of Physics, The University of Texas at Austin, Austin, Texas 78712-1081

(Received 30 March 2001; published 10 July 2001)

We report the first observation of the quantum Zeno and anti-Zeno effects in an unstable system. Cold sodium atoms are trapped in a far-detuned standing wave of light that is accelerated for a controlled duration. For a large acceleration the atoms can escape the trapping potential via tunneling. Initially the number of trapped atoms shows strong nonexponential decay features, evolving into the characteristic exponential decay behavior. We repeatedly measure the number of atoms remaining trapped during the initial period of nonexponential decay. Depending on the frequency of measurements we observe a decay that is suppressed or enhanced as compared to the unperturbed system.




FIG. 3. Probability of survival in the accelerated potential as a function of duration of the tunneling acceleration. The hollow squares show the noninterrupted sequence, and the solid circles show the sequence with interruptions of $50 \mu\text{s}$ duration every $1 \mu\text{s}$. The error bars denote the error of the mean. The data have been normalized to unity at $t_{\text{tunnel}} = 0$ in order to compare with the simulations. The solid lines are quantum mechanical simulations of the experimental sequence with no adjustable parameters. For these data the parameters were $a_{\text{tunnel}} = 15\,000 \text{ m/s}^2$, $a_{\text{interr}} = 2000 \text{ m/s}^2$, $t_{\text{interr}} = 50 \mu\text{s}$, and $V_0/h = 91 \text{ kHz}$, where h is Planck's constant.

Zeno effekt

Same exp. setup,
but with measurements in between




FIG. 4. Survival probability as a function of duration of the tunneling acceleration. The hollow squares show the noninterrupted sequence, and the solid circles show the sequence with interruptions of $40 \mu\text{s}$ duration every $5 \mu\text{s}$. The error bars denote the error of the mean. The experimental data points have been connected by solid lines for clarity. For these data the parameters were: $a_{\text{tunnel}} = 15\,000 \text{ m/s}^2$, $a_{\text{interr}} = 2800 \text{ m/s}^2$, $t_{\text{interr}} = 40 \mu\text{s}$, and $V_0/h = 116 \text{ kHz}$.

Anti-Zeno effect

Late-time deviations

PRL 96, 163601 (2006)

PHYSICAL REVIEW LETTERS

week ending
28 APRIL 2006



Violation of the Exponential-Decay Law at Long Times

C. Rothe, S. I. Hintschich, and A. P. Monkman

Department of Physics, University of Durham, Durham, DH1 3LE, United Kingdom

(Received 4 July 2005; published 26 April 2006)

First-principles quantum mechanical calculations show that the exponential-decay law for any metastable state is only an approximation and predict an asymptotically algebraic contribution to the decay for sufficiently long times. In this Letter, we measure the luminescence decays of many dissolved organic materials after pulsed laser excitation over more than 20 lifetimes and obtain the first experimental proof of the turnover into the nonexponential decay regime. As theoretically expected, the strength of the nonexponential contributions scales with the energetic width of the excited state density distribution whereas the slope indicates the broadening mechanism.




FIG. 2 (color). Corresponding double logarithmic fluorescence decays of the emissions shown in Fig. 1. Exponential and power law regions are indicated by solid lines and the emission intensity at time zero has been normalized.

Confirmation of: L. A. Khal'fin. 1957. 1957 (Engl. trans. Zh.Eksp.Teor.Fiz.,33,1371)

Francesco Giacosa

Considerations



- No other short- or long-time deviation from the exp. law was seen in unstable states.
- Verification of the two aforementioned works (Reizen + Rothe) would be needed.
- The measurement of deviations in simple natural systems (elementary particles, nuclei, atoms) would be a great achievement.

Part 3: neutron decay anomaly

Neutron decay: exp. methods



- There are two methods to measure the lifetime of neutrons: beam and trap
- Beam: one measures the protons out of a neutron beam
- Trap: one measures the neutrons that survive in a certain neutron trap

Exp. results: beam vs trap/bottle

$$\tau_n^{\text{beam}} = 888.1 \pm 2.0 \text{ s}$$

$$\Delta\tau = \tau_n^{\text{beam}} - \tau_n^{\text{trap}} = 8.7 \pm 2.1 \text{ s}$$


$$\tau_n^{\text{trap}} = 879.45 \pm 0.58 \text{ s}$$

$$\Gamma_n^{\text{trap}} / \Gamma_n^{\text{beam}} = 1.0098 \pm 0.0024$$

Article

Measurements of the Neutron Lifetime

F. E. Wietfeldt



Hidden decays of the neutron?



the proposal of Fornal and Grinstein

B. Fornal and B. Grinstein, Phys. Rev. Lett. **120**, 191801 (2018), 1801.01124.

The beam experiments shows a larger lifetime because it misses some of (BSM) decays of the neutron. In particular, decays into a light fermion n ;

However, the existence of such a light fermion is at odds with bounds from neutron stars, see:

D. McKeen, A. E. Nelson, S. Reddy, and D. Zhou, Phys. Rev. Lett. **121**, 061802 (2018), 1802.08244.

G. Baym, D. H. Beck, P. Geltenbort, and J. Shelton, Phys. Rev. Lett. **121**, 061801 (2018), 1802.08282.

T. F. Motta, P. A. M. Guichon, and A. W. Thomas, J. Phys. **G45**, 05LT01 (2018), 1802.08427.



Part 4: IZE and neutron decay

1906.10024

Neutron decay anomaly and inverse quantum Zeno effect

Francesco Giacosa^{(a,b)*} and Giuseppe Pagliara^{(c)†}

IZE as a possible explanation of the neutron decay anomaly



A. Kofman and G. Kurizki, *Zeitschrift für Naturforschung A* **56** (2001).
Francesco Giacosa

Environment and effective decay width

$$\Gamma_n^{\text{measured}}(\tau) = \int_0^\infty f(\tau, \omega) \Gamma_n(\omega) d\omega$$

$\Gamma(\omega)$ is the decay width, $f(\tau, \omega)$ the response of the environment

$$\int_0^\infty f(\tau, \omega) d\omega = 1$$

$$f(\tau \rightarrow \infty, \omega) = \delta(\omega - \omega_n)$$

$$\Gamma_n^{\text{measured}}(\tau \rightarrow \infty) = \Gamma_n^{\text{on-shell}}$$

$$f(\tau \rightarrow 0, \omega) = \text{small const}$$

A. Kofman and G. Kurizki, Nature **405**, 546 (2000).

A. Kofman and G. Kurizki, Zeitschrift für Naturforschung A **56** (2001).

P. Facchi and S. Pascazio, Fortschritte der Physik **49**, 941 (2001).

Two explicit form of the response function $f(\tau, \omega)$



$$f(\tau, \omega) = \frac{\tau}{2\pi} \frac{\sin^2[(\omega - \omega_n)/2]}{[(\omega - \omega_n)/2]^2}$$

This is for ideal collapses at $\tau, 2\tau, 3\tau, \dots$

$$f(\tau, \omega) = \left[(\omega - \omega_n)^2 + \tau^{-2} \right]^{-1} / \pi\tau$$

This is for a continuous measurement

The case of the neutron

$$\Gamma_n(\omega) = g_n^2 \omega^5 \quad \text{for } \omega \lesssim \omega_{\text{on-shell}} + m_\pi$$


$$g_n \propto g_V V_{ud}$$

$$\Gamma_n^{\text{on-shell}} = \Gamma_n(\omega_n) = g_n^2 \omega_n^5 = \hbar/888.1 \text{ sec}^{-1} = 7.41146 \cdot 10^{-25} \text{ MeV}$$

$$\omega_n = \omega_{\text{on-shell}} = m_n^{\text{on-shell}} - m_p - m_e = 0.782333 \text{ MeV}$$

Note: for very large ω the decay width function should go to zero

The basic idea



Integral over $\Gamma(\omega)$ and $f(\tau, \omega)$

**QUANTUM ZENO EFFECTS WITH “PULSED”
AND “CONTINUOUS” MEASUREMENTS**

P. Facchi⁽¹⁾ and S. Pascazio⁽²⁾

INTERNATIONAL CONFERENCE
TAQMSB

**TIME'S ARROWS, QUANTUM MEASUREMENT
AND SUPERLUMINAL BEHAVIOR**

Introducing an upper 'energy' for convergence

$$\Gamma_n^{\text{measured}}(\tau, \omega_C) = \int_0^{\omega_C} f(\tau, \omega) \Gamma_n(\omega) d\omega$$

ω_C measures the maximal off-shellness of the neutron;
It should be of the order of 1-10 MeV.

$$\Gamma_n^{\text{measured}}(\tau, \omega_C) = \Gamma_n^{\text{on-shell}} \left(1 + \frac{\hbar}{\tau} \frac{\omega_C^4}{4\pi\omega_n^5} \right)$$

$$\Gamma_n^{\text{measured}}(\tau, \omega_C) > \Gamma_n^{\text{on-shell}}$$

Discussion



- Beam: how often is the decay determined? Not that often, $\tau=10^{-9}$ s.
- Traps: Here, we have typically a set of 10^8 cold neutrons entangled in a Slater determinant. Measuring one means to collapse them all. $\tau=10^{-17}$ s.

IZE: Beam vs Trap

$$\Gamma_n^{\text{beam}} \simeq \Gamma_n^{\text{measured}}(\tau \sim 10^{-9}, \omega_C \sim 2-10\omega_n) \simeq \Gamma_n^{\text{on-shell}}$$

For the beam: exponential on-shell result

$$\Gamma_n^{\text{trap}} \simeq \Gamma_n^{\text{measured}}(\tau \sim 10^{-17}, \omega_C \sim 2-10\omega_n) \gtrsim \Gamma_n^{\text{on-shell}}$$

For the trap: possible increase of Γ


$$\tau \sim 10^{-9} \cdot 10^{-8} = 10^{-17} \text{ s.}$$

$$\omega_C = 6.19\omega_n$$

$$\Gamma_n^{\text{trap}} = 1.0098\Gamma_n^{\text{on-shell}} = 1.0098\Gamma_n^{\text{beam}}$$

See 1906.10024 for details

ω_C vs T



1906.10024

Conclusions



- QZE and IZE are a well-established part of QM
- The IZE has been presented as a possible solution of the neutron decay anomaly
- Increasing/decreasing of nr of neutrons in the trap may have an influence on the measured values.
- Measuring the protons in trap exps should confirm the smaller decay width (no influence)

Thank You

Lee Hamiltonian



$$H = H_0 + H_1$$

$$H_0 = M_0 |S\rangle\langle S| + \int_{-\infty}^{+\infty} dk \omega(k) |k\rangle\langle k|$$

$$H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g \cdot f(k)) (|S\rangle\langle k| + |k\rangle\langle S|)$$

$|S\rangle$ is the initial unstable state, coupled to an infinity of final states $|k\rangle$. (Poincare-time is infinite. Irreversible decay). General approach, similar Hamiltonians used in many areas of Physics.

(Ex: Jaynes-Cummings approach)

Example/1: spontaneous emission. $|S\rangle$ represents an atom in the excited state, $|k\rangle$ is the ground-state plus photon.

Example/2: pion decay. $|S\rangle$ represents a neutral pion, $|k\rangle$ represents two photons (flying back-to-back)


Exponential limit

$$H = H_0 + H_1 ; H_0 = M_0 |S\rangle\langle S| + \int_{-\infty}^{+\infty} dk \omega(k) |k\rangle\langle k| ; H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g \cdot f(k)) (|S\rangle\langle k| + |k\rangle\langle S|)$$

$$\omega(k) = k ; f(k) = 1 \Rightarrow \Pi(E) = ig^2 / 2 ; \Gamma = g^2$$

$$d_s(E) = \frac{\Gamma}{2\pi} \frac{1}{(E - M_0)^2 + \Gamma^2 / 4}$$

$$\Rightarrow a(t) = e^{-i(M_0 - i\Gamma/2)t} \Rightarrow p(t) = e^{-\Gamma t}$$




The exponential limit is obtained when the unstable state couples to all the states of the continuum with the same strength

Non-exponential case (1)

$$H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g \cdot f(k)) (|S\rangle\langle k| + |k\rangle\langle S|)$$


$$f(k) = \begin{cases} 0 & \text{for } k < E_{\min} \\ 1 & \text{for } E_{\min} \leq k \leq E_{\max} \\ 0 & \text{for } k > E_{\max} \end{cases}$$




$$M_0 = 2; E_{\min} = 0; E_{\max} = 5; g^2 = 0.36 \text{ (all in a.u. of energy)}$$

This is what I have said at the beginning of the talk, but now “well done”

Non-exponential case (2)



Dashed: $p_{\text{BW}}(t) = e^{-\Gamma t}$ with $\Gamma = \text{Im}[\Pi(M)] / 2$



$$h(t) = -\frac{dp(t)}{dt}$$

Dashed: $h_{\text{BW}}(t) = \Gamma e^{-\Gamma t}$ with $\Gamma = \text{Im}[\Pi(M)] / 2$

$$\int_0^t h(u) du = 1 - p(t)$$

Namley: $h(t)dt = p(t) - p(t + dt)$ is the probability that the particles decays between t and $t + dt$

Two-channel case (a)

Found Phys (2012) 42:1262–1299
DOI 10.1007/s10701-012-9667-3




Non-exponential Decay in Quantum Field Theory and in Quantum Mechanics: The Case of Two (or More) Decay Channels

Francesco Giacosa

$$H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g_1 \cdot f_1(k)) (|S\rangle \langle k, 1| + |k, 1\rangle \langle S|) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g_2 \cdot f_2(k)) (|S\rangle \langle k, 2| + |k, 2\rangle \langle S|)$$

$$f_i(k) = \begin{cases} 0 & \text{for } k < E_{i,\min} \\ 1 & \text{for } E_{i,\min} \leq k \leq E_{i,\max} \\ 0 & \text{for } k > E_{i,\max} \end{cases}$$



$$M_0 = 2; E_{1,\min} = 0; E_{2,\min} = 0; E_{1,\max} = E_{2,\max} = 5;$$


$$g_1^2 = 0.36; g_2^2 = 0.16 \quad (\text{all in a.u. of energy})$$

Francesco Giacosa

Two-channel case (b)

$h_1(t)dt =$ probability that the state $|S\rangle$ decays in the first channel between $(t,t+dt)$

$h_2(t)dt =$ probability that the state $|S\rangle$ decays in the second channel between $(t,t+dt)$



Measurable effect???

Details in:

F. G., Non-exponential decay in quantum field theory and in quantum mechanics: the case of two (or more) decay channels, Found. Phys. 42 (2012) 1262 [arXiv:1110.5923].

Experimental confirmation of the quantum Zeno effect - Itano et al (1)

PHYSICAL REVIEW A

VOLUME 41, NUMBER 5

1 MARCH 1990


Quantum Zeno effect

Wayne M. Itano, D. J. Heinzen, J. J. Bollinger, and D. J. Wineland

Time and Frequency Division, National Institute of Standards and Technology, Boulder, Colorado 80303

(Received 12 October 1989)

The quantum Zeno effect is the inhibition of transitions between quantum states by frequent measurements of the state. The inhibition arises because the measurement causes a collapse (reduction) of the wave function. If the time between measurements is short enough, the wave function usually collapses back to the initial state. We have observed this effect in an rf transition between two ${}^9\text{Be}^+$ ground-state hyperfine levels. The ions were confined in a Penning trap and laser cooled. Short pulses of light, applied at the same time as the rf field, made the measurements. If an ion was in one state, it scattered a few photons; if it was in the other, it scattered no photons. In the latter case the wave-function collapse was due to a null measurement. Good agreement was found with calculations.



(Undisturbed) survival probability

At $t = 0$, the electron is in $|1\rangle$.

$$p(t) = \cos^2\left(\frac{\Omega t}{2}\right) = 1 - \frac{\Omega^2 t^2}{4} + \dots$$

$$p(T) = 0 \text{ für } T = \pi/\Omega$$

Experimental confirmation of the quantum Zeno effect - Itano et al (2)





FIG. 2. Diagram of the energy levels of ${}^9\text{Be}^+$ in a magnetic field B . The states labeled 1, 2, and 3 correspond to those in Fig. 1.

5000 ions in a Penning trap

Short laser pulses 1-3 work as measurements.




FIG. 3. Graph of the experimental and calculated $1 \rightarrow 2$ transition probabilities as a function of the number of measurement pulses n . The decrease of the transition probabilities with increasing n demonstrates the quantum Zeno effect.

$$p(t) = \cos^2(\Omega t / 2) = 1 - \frac{\Omega^2 t^2}{4} + \dots ; \quad p(T) = 0 \text{ für } T = \pi/\Omega$$

(Transition probability (without measuring) at time T): $1 - p(T) = 1$.

With n measurements in between the transition probability decreases!

The electron stays in state 1.

Other experiments about Zeno/Streed et al

PRL **97**, 260402 (2006)

PHYSICAL REVIEW LETTERS

week ending
31 DECEMBER 2006

Continuous and Pulsed Quantum Zeno Effect

Erik W. Streed,^{1,2} Jongchul Mun,¹ Micah Boyd,¹ Gretchen K. Campbell,¹ Patrick Medley,¹
Wolfgang Ketterle,¹ and David E. Pritchard¹

¹*Department of Physics, MIT-Harvard Center for Ultracold Atoms, and Research Laboratory of Electronics, MIT,
Cambridge, Massachusetts 02139, USA*

²*Centre for Quantum Dynamics, Griffith University, Nathan, QLD 4111, Australia*
(Received 14 June 2006; published 27 December 2006)

Use of BEC (with Rb). QZE confirmed.

The intensity of a continuous observation of a quantum state is equivalent to a certain t_0 (Shulman, PRA 57, 1509 (1997)).

Other experiments about Zeno/Haroche



PRL **101**, 180402 (2008)

PHYSICAL REVIEW LETTERS

week ending
31 OCTOBER 2008



Freezing Coherent Field Growth in a Cavity by the Quantum Zeno Effect

J. Bernu,¹ S. Deléglise,¹ C. Sayrin,¹ S. Kuhr,^{1,*} I. Dotsenko,^{1,2} M. Brune,^{1,+} J. M. Raimond,¹ and S. Haroche^{1,2}

¹*Laboratoire Kastler Brossel, Ecole Normale Supérieure, CNRS, Université P. et M. Curie,
24 rue Lhomond, F-75231 Paris Cedex 05, France*

²*Collège de France, 11 Place Marcelin Berthelot, F-75231 Paris Cedex 05, France*



(Received 21 July 2008; published 28 October 2008)

Cavity QED: the nr of photons is frozen.

Another verification of QZE.

Direction QFT.

Other experiments about Zeno/Balzer



Same setup as Itano et al. (different ions are used, YB instead of Be),

But now the measurement takes place between 3 and 2.

Results in agreement with Itano, but here the QZE is associated by a series of null-measurements.

Quantum Zeno dynamics, Quantum computations, ...



Sudarshan: Seven Science Quests

IOP Publishing

Journal of Physics: Conference Series 196 (2009) 012017

doi:10.1088/1742-6596/196/1/012017

Quantum Zeno dynamics and quantum Zeno subspaces

Paolo Facchi¹, Giuseppe Marmo², Saverio Pascazio³

¹ Dipartimento di Matematica, Università di Bari
and Istituto Nazionale di Fisica Nucleare, Sezione di Bari, I-70125 Bari, Italy

² Dipartimento di Scienze Fisiche, Università di Napoli "Federico II"

and Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, I-80126 Napoli, Italy

³ Dipartimento di Fisica, Università di Bari

and Istituto Nazionale di Fisica Nucleare, Sezione di Bari, I-70126 Bari, Italy

E-mail: saverio.pascazio@ba.infn.it

PRL 108, 080501 (2012)

PHYSICAL REVIEW LETTERS

week ending
24 FEBRUARY 2012

Zeno Effect for Quantum Computation and Control

Gerardo A. Paz-Silva,^{1,4} A. T. Rezakhani,^{1,4,5} Jason M. Dominy,^{1,4} and D. A. Lidar^{1,2,3,4}

¹Department of Chemistry, University of Southern California, Los Angeles, California 90089, USA

²Department of Electrical Engineering, University of Southern California, Los Angeles, California 90089, USA

³Department of Physics, University of Southern California, Los Angeles, California 90089, USA

⁴Center for Quantum Information Science and Technology, University of Southern California, Los Angeles, California 90089, USA

⁵Department of Physics, Sharif University of Technology, Tehran, Iran

(Received 28 April 2011; revised manuscript received 9 February 2012; published 22 February 2012)

C. R. Physique 17 (2016) 685–692



Contents lists available at ScienceDirect

Comptes Rendus Physique

www.sciencedirect.com



Quantum microwaves / Micro-ondes quantiques

Quantum Zeno dynamics in atoms and cavities

Dynamique de Zénon quantique avec des atomes et des cavités

Sébastien Gleyzes*, Jean-Michel Raimond

Laboratoire Kastler Brossel, Collège de France, CNRS, ENS-PSL Research University, UPMC-Sorbonne Universités, 11, place Marcelin-Berthelot, 75005 Paris, France



Francesco Giacosa