3rd Jagiellonian Symposium on Fundamental and Applied Subatomic Physics



Contribution ID: 72 Type: poster

Response functions and cross sections for inclusive neutrino scattering off ²H

Tuesday, 25 June 2019 13:30 (1h 30m)

Neutrino scattering on nuclei has been investigated for several decades. It played an important role in establishing fundamentals of the theory of weak interactions and electroweak unification. Later it became a tool to study the much more subtle properties of neutrinos, such as masses and oscillations.

Most of the calculations for neutrino scattering on light nuclei were performed in coordinate space, see important Refs. (1-4). In a recent paper (5) we presented momentum-space based results for several (anti)neutrino induced reactions on light nuclei: $\nu(\bar{\nu}) + {}^2{\rm H} \rightarrow \nu(\bar{\nu}) + p + n$,

```
\begin{split} \bar{\nu} &+ {}^{2}\mathrm{H} \rightarrow e^{+} + n + n\,, \\ \nu(\bar{\nu}) &+ {}^{3}\mathrm{He} \rightarrow \nu(\bar{\nu}) + p + {}^{2}\mathrm{H}\,, \\ \nu(\bar{\nu}) &+ {}^{3}\mathrm{He} \rightarrow \nu(\bar{\nu}) + p + p + n\,, \\ \nu(\bar{\nu}) &+ {}^{3}\mathrm{H} \rightarrow \nu(\bar{\nu}) + p + {}^{2}\mathrm{H}\,, \\ \nu(\bar{\nu}) &+ {}^{3}\mathrm{H} \rightarrow \nu(\bar{\nu}) + n + n + p\,, \\ \nu &+ {}^{3}\mathrm{He} \rightarrow e^{+} + n + {}^{2}\mathrm{H}\,, \\ \nu &+ {}^{3}\mathrm{He} \rightarrow e^{+} + n + n + p\,, \end{split}
```

 $\nu+^3{
m H} o e^++n+n+n$. We restricted our selves to neutrino energie supto 300 MeV and pointed out to problems arising from mergical many incoming neutrino energies. The essential building block for the total cross section at a given (anti)neutrino energy E is the threefold differential cross section $d^3\sigma/d\Omega'/dE'$, which depends on E, on the (anti)neutrino scattering angle θ' and on its final energy E'.

Namely, the total cross section is most easily calculated as $\sigma_{tot}(E) = \int d\Omega' \int dE' \frac{d^3\sigma}{d\Omega' dE'} = 2\pi \int d\theta' \sin \theta' \int dE' \frac{d^3\sigma}{d\Omega' dE'}$. The cross can be expressed in terms of the so-called response functions R_{jk} (5), which depend on two parameters only: on the internal energy of the nuclear system E_{CM} and on the magnitude of the three-momentum transfer Q: $\mathrm{d}^3\sigma \frac{\partial G'}{\partial G'} \frac{\partial G$

are analytically known, it is thus much more efficient to calculate just the response functions on a sufficiently dense two dimensional grid in the (E_{CM},Q) plane and use interpolations to obtain these values of the response functions which are needed to compute $\sigma_{tot}(E)$ at a required energy E. Note also that in the case of the neutral current driven reactions, exactly the same response functions can be used to obtain cross sections with neutrinos and antineutrinos. This approach has been already successfully tested for the (anti)neutrino reactions on the deuteron and one example of the response function for the $\nu(\bar{\nu}) + {}^2{\rm H} \rightarrow \nu(\bar{\nu}) + p + n$ reaction is shown in Fig. 1. The corresponding work for several reactions with the three-nucleon systems is in progress and will be reported at the conference. As in Ref. (5), also in the present calculations the single nucleon current from Ref. (6) and the AV18 two-nucleon potential (7) is employed.

References

(1) S. Nakamura, T. Sato, V. Gudkov, and K. Kubodera, Phys. Rev. C 63, 034617 (2001);

Erratum: [Phys. Rev. C 73, 049904 (2006)].

(2) S. Nakamura et al., Nucl. Phys. A 707, 561 (2002);

http://www-nuclth.phys.sci.osaka-u.ac.jp/top/Netal/total/index.html.

- (3) G. Shen, L. E. Marcucci, J. Carlson, S. Gandolfi, and R. Schiavilla, Phys. Rev. C 86, 035503 (2012).
- (4) A. Baroni and R. Schiavilla, Phys. Rev. C 96, 014002 (2017).
- (5) J. Golak et al., Phys. Rev. C 98, 015501 (2018).

- (6) J. Golak et al., Phys. Rev. C 90, 024001 (2014).
- (7) R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, Phys. Rev. C ${\bf 51}$, 38 (1995).

Primary author: GRASSI, Alessandro ((for LENPIC Collaboration))

Presenter: GRASSI, Alessandro ((for LENPIC Collaboration))

Session Classification: Poster session