

Search for eta-mesic helium with WASA-at-COSY*

MAGDALENA SKURZOK^{a,b}
FOR THE WASA-AT-COSY COLLABORATION

^aINFN, Laboratori Nazionali di Frascati, 00044 Frascati, Italy

^bM. Smoluchowski Institute of Physics, Jagiellonian University, Cracow, Poland

We report on the search for $^4\text{He-}\eta$ and $^3\text{He-}\eta$ mesic nuclei with WASA-at-COSY detection system. The description of the experimental method as well as recent status of the data analysis are presented.

PACS numbers: 1.85.+d, 21.65.Jk, 25.80.-e, 13.75.-n

1. Introduction

The mesic nuclei, an exotic nuclear matter consisting of a nucleus bound via the strong interaction with a neutral meson (η , η' , K or ω), are currently one of the hottest topic in nuclear and hadronic physics, both from experimental [1–7] and theoretical side [9–32].

η meson is one of the most promising candidates for the creation of the mesic nucleus since its interaction with nucleons is stronger than other mesons [33, 34]. Current investigations of the η meson production result in a wide range of ηN scattering length values ($a_{\eta N}$) indicating the η -nucleon interaction to be strong enough to create light η -mesic bound states [11–14, 35–37]. However, none of performed experiments have brought clear evidence of their existence [38–45]. Recent reviews concerning η mesic nuclei searches one can find in Refs [6–8, 19, 21, 33, 46–50].

Promising experiments related to η -mesic Helium nuclei have been performed recently with the WASA-at-COSY facility. The search for $^4\text{He-}\eta$ and $^3\text{He-}\eta$ mesic bound systems has been carried out in dd and pd collisions, respectively, using unique ramped beam technique. This paper reports on the results obtained for the search for η -mesic ^4He in $dd \rightarrow ^3\text{He}n\pi^0$ and

* Presented at 3rd Jagiellonian Symposium on Fundamental and Applied Subatomic Physics

$dd \rightarrow {}^3\text{He}p\pi^-$ processes [1–4, 51]. Preliminary results for ${}^3\text{He}\eta$ mesic nuclei searches in $pd \rightarrow {}^3\text{He}2\gamma(6\gamma)$ reactions are also presented [53, 54].

2. Search for the η -mesic ${}^4\text{He}$

WASA-at-COSY Collaboration performed two measurements dedicated to search for η -mesic ${}^4\text{He}$ nuclei (in 2008 and 2010) using the unique ramped beam technique which allows for the slow and continuous beam momentum changes around the η production threshold in each of the beam acceleration cycle [1, 3, 48, 50, 51]. The uniqueness of this technique lies in the reduction of systematic uncertainties with respect to separate runs at fixed beam energies [3, 41].

In order to search for ${}^4\text{He}\eta$ bound states the excitation functions for $dd \rightarrow {}^3\text{He}p\pi^-$ [1–4] and $dd \rightarrow {}^3\text{He}n\pi^0$ [1, 2, 4] reactions have been studied around the ${}^4\text{He}\eta$ production threshold ($Q \in (-70, 30)$ MeV). A detailed description of the data analysis is presented in Refs [1, 3]. Since the obtained excitation functions do not reveal any direct signature of the bound state below the η production threshold, the upper limit of the total cross section for the η -mesic ${}^4\text{He}$ formation and its decay into each of investigated channel was determined at the 90% confidence level. Therefore, the excitation curves were fitted simultaneously using the function being a sum of Breit-Wigner function with a fixed binding energy and width and a second order polynomial for signal and background, respectively. During the fit the isospin relation between $n\pi^0$ and $p\pi^-$ pairs has been taken into account.

The upper limit of the total cross section for $dd \rightarrow ({}^4\text{He}\eta)_{\text{bound}} \rightarrow {}^3\text{He}n\pi^0$ and $dd \rightarrow ({}^4\text{He}\eta)_{\text{bound}} \rightarrow {}^3\text{He}p\pi^-$ processes varies in the range from 2.5 to 3.5 nb and 5 to 7 nb, respectively. In case of the first process the upper limit was determined experimentally for the first time, while the sensitivity of the cross section achieved for the second reaction was about four times better in comparison with the result obtained in the previous experiment [3]. The obtained upper limits as a function of the bound state width are presented for both studied processes in Fig. 1.

The data analysis presented above was carried out under assumption that the signal from the bound state is described by a Breit-Wigner shape with fixed binding energy and width [1, 3]. However, a theoretical description of the cross sections in the excess energy range relevant to the η -mesic nuclear search was proposed in Ref. [10]. A phenomenological approach with an optical potential for the η - ${}^4\text{He}$ interaction was applied and the total cross sections were determined for a broad range of real (V_0) and imaginary (W_0) parameters.

The upper limit of the total cross section (CL=90%) for creation of η -




Fig. 1. Upper limit of the total cross-section for $dd \rightarrow (^4\text{He}-\eta)_{\text{bound}} \rightarrow {}^3\text{He} n \pi^0$ (left panel) and $dd \rightarrow (^4\text{He}-\eta)_{\text{bound}} \rightarrow {}^3\text{He} p \pi^-$ (right panel) reaction as a function of the width of the bound state. The binding energy was fixed to 30 MeV. The upper limit was determined via the simultaneous fit for both channels. The green area denotes the systematic uncertainties. The figures are adopted from [1].

mesic nuclei via the $dd \rightarrow {}^3\text{He} N \pi$ was determined by fitting the theoretical spectra convoluted with the experimental resolution of the excess energy to experimental data collected by WASA-at-COSY [1]. It was found to vary from about 5.2 nb to about 7.5 nb [2].

Comparing the determined upper limits with the cross sections obtained in Ref. [10] we were able to put a constraint on the η - ${}^4\text{He}$ optical potential parameters. As it is presented in Fig. 2, most optical model predictions [10] are excluded except for extremely narrow and loosely bound states. Details of performed studies are presented in Ref. [2].

3. Search for the η -mesic ${}^3\text{He}$

A promising experiment dedicated to the search for η -mesic ${}^3\text{He}$ was performed with WASA-at-COSY facility using ramped proton beam ($Q \in (-70, 30)$ MeV) and deuteron target. Three different mechanisms of the η -mesic bound state decay were considered: (i) the η meson absorption by one of the nucleons leading to excitation of $N^*(1535)$ resonance which subsequently decays into nucleon-pion pair (assumed as well in data analysis and interpretations of previous experiments [1–3]), (ii) η -meson decay while it is still “orbiting” around a nucleus and (iii) two-nucleon η meson absorption process.

The theoretical model (ii) was developed recently [52] and has been applied in analysis of $pd \rightarrow ({}^3\text{He}-\eta)_{\text{bound}} \rightarrow {}^3\text{He} 2\gamma$ and $pd \rightarrow ({}^3\text{He}-\eta)_{\text{bound}} \rightarrow {}^3\text{He} 6\gamma$ decay channels [53]. Excitation functions determined for both chan-




Fig. 2. Contour plot of the theoretically determined conversion cross section in the $V_0 - W_0$ plane [10]. The light shaded area shows the region excluded by our analysis, while the dark shaded area denotes the systematic uncertainty of the $\sigma_{upper}^{CL=90\%}$. The red line extends the allowed region based on a new estimate of errors (see text for details). Dots correspond to the optical potential parameters corresponding to the predicted η -mesic ${}^4\text{He}$ states. Figure is adopted from Ref. [2].

nels do not show the bound state signature, therefore, the upper limit of the total cross section at the CL=90% was determined for the η -mesic ${}^3\text{He}$ nucleus creation followed by the η meson decay. For this purpose excitation functions for both reactions were simultaneously fitted with a Breit-Wigner function (signal) combined with polynomial (background). During the analysis the branching ratio relation between $\eta \rightarrow 2\gamma$ and $\eta \rightarrow 3\pi^0$ in vacuum was taken into account. The preliminary estimated upper limit varies between 2 nb to 15 nb depending on the bound state parameters (binding energy, width) [53].

4. Summary and perspectives

Search for η -mesic Helium was performed with the WASA-at-COSY facility in deuteron-deuteron and proton-deuteron collisions. Resonance structure related to the η -mesic ${}^4\text{He}$ bound state was not observed in $dd \rightarrow {}^3\text{He}n\pi^0$ and $dd \rightarrow {}^3\text{He}p\pi^-$ reactions. However, the upper limits of the total cross sections for η -mesic nuclei formation and decay in each channel were determined to be of the order of a few nb [1, 3]. In addition, the experimental data has been compared with the phenomenological model proposed in Ref. [10] allowing, for the first time, to constrain the range of the η - ${}^4\text{He}$ optical potential parameters [2].

In 2014, an experiment dedicated to the search for η -mesic ${}^3\text{He}$ in three

different mechanisms has been performed. The measurement with a high average luminosity ($\sim 3 \cdot 10^{30} \text{ cm}^{-2} \text{ s}^{-1}$) allowed the collection of the largest data sample, in the world available up to now, for ${}^3\text{He}-\eta$ [54–56]. The preliminary upper limit value for $pd \rightarrow {}^3\text{He}2\gamma$ and $pd \rightarrow {}^3\text{He}6\gamma$ channels is on the level of a few nanobarns. The analysis assuming a mechanism of bound state decay via N^* excitation and its decay into the nucleon-pion pair is in progress [57].

REFERENCES

- [1] P. Adlarson *et al.*, Nucl. Phys. A **959**, 102-115 (2017)
- [2] M. Skurzok *et al.*, Phys. Lett. B **772**, 663 (2018)
- [3] P. Adlarson *et al.*, Phys. Rev. C **87**, 035204 (2013)
- [4] M. Skurzok, P. Moskal, W. Krzemien, Prog. Part. Nucl. Phys. **67**, 445 (2012)
- [5] Y. K. Tanaka *et al.*, Phys. Rev. Lett. **117**, 202501 (2016)
- [6] H. Machner, J. Phys. G **42**, 043001 (2015)
- [7] V. Metag, M. Nanova and E. Ya. Paryev, Prog. Part. Nucl. Phys. **97**, 199 (2017)
- [8] S. D. Bass, P. Moskal, Rev. Mod. Phys. **91**, 015003 (2019)
- [9] J.-J. Xie *et al.*, Eur. Phys. J. A **55** no.1, 6 (2019)
- [10] N. Ikeno *et al.*, Eur. Phys. J. A **53** no.10, 194 (2017)
- [11] J. J. Xie *et al.*, Phys. Rev. C **95**, 015202 (2017)
- [12] A. Fix *et al.*, Phys. Lett. B **772**, 663 (2017)
- [13] N. Barnea, B. Bazak, E. Friedman and A. Gal, Phys. Lett. B **771**, 297 (2017)
- [14] N. Barnea, E. Friedman and A. Gal, Nucl. Phys. A **968**, 35 (2017)
- [15] T. Sekihara, H. Fujioka, T. Ishikawa, Phys. Rev. C **97**, 045202 (2017)
- [16] N. Barnea, E. Friedman and A. Gal, Phys. Lett. B **747**, 345 (2015)
- [17] E. Friedman, A. Gal, J. Mares, Phys. Lett. B **725**, 334 (2013)
- [18] N. G. Kelkar, Eur. Phys. J. A **52**, 309 (2016)
- [19] N. G. Kelkar *et al.*, Rept. Progr. Phys. **76**, 066301 (2013)
- [20] N. G. Kelkar, Acta Phys. Pol. B **46**, 113 1 (2015)
- [21] C. Wilkin, Acta Phys. Pol. B **47**, 249 (2016)
- [22] C. Wilkin, Phys. Lett. B **654**, 92 (2007)
- [23] S. D. Bass, A. W. Thomas, Phys. Lett. B **634**, 368 (2006)
- [24] S. D. Bass, A. W. Thomas, Acta Phys. Polon. B **45**, 627 (2014)
- [25] S. Hirenzaki and H. Nagahiro, Acta Phys. Polon. B **45**, 619 (2014)
- [26] H. Nagahiro, D. Jido and S. Hirenzaki, Phys. Rev. C **80**, 025205 (2009)

- [27] H. Nagahiro *et al.*, Phys. Rev. C **87**, 045201 (2013)
- [28] S. Hirenzaki *et al.*, Acta Phys. Polon. B **41**, 2211 (2010)
- [29] S. Wycech, W. Krzemien, Acta Phys. Pol. B **45**, 745 (2014)
- [30] J. Niskanen, Phys. Rev. C **92**, 055205 (2015)
- [31] Q. Haider, L. C. Liu, Phys. Lett. B **172**, 257 (1986)
- [32] R. S. Bhalariao and L. C. Liu, Phys. Rev. Lett. **54**, 865 (1985)
- [33] P. Moskal, Few Body Syst. **50**, 667 (2014)
- [34] P. Moskal *et al.*, Phys. Lett. B **482**, 356 (2000)
- [35] A. M. Green, J. A. Niskanen, S. Wycech, Phys. Rev. C **54**, 1970 (1996)
- [36] C. Wilkin, Phys. Rev. C **47**, 938 (1993)
- [37] S. Wycech, A. M. Green and J. A. Niskanen, Phys. Rev. C **52**, 544 (1995)
- [38] J. Berger *et al.*, Phys. Rev. Lett. **61**, 919 (1988)
- [39] B. Mayer *et al.*, Phys. Rev. C **53**, 2068 (1996)
- [40] G. A. Sokol, L. N. Pavlyuchenko, *arXiv:nucl-ex/0111020* (2001)
- [41] J. Smyrski *et al.*, Phys. Lett. B **649**, 258 (2007)
- [42] T. Mersmann *et al.*, Phys. Rev. Lett. **98**, 242301 (2007)
- [43] A. Budzanowski, *et al.*, Phys. Rev. C **79**, 012201(R) (2009)
- [44] M. Papenbrock, *et al.*, Phys. Lett. B **734**, 333 (2014)
- [45] P. Moskal, J. Smyrski, Acta Phys. Polon. B **41**, 2281 (2010)
- [46] Q. Haider, L.-C. Liu, J. Phys. G **37**, 125104 (2010)
- [47] B. Krusche, C. Wilkin, Prog. Part. Nucl. Phys. **80**, 43 (2014)
- [48] M. Skurzok *et al.*, Acta Phys. Polon. B **47**, 503 (2016)
- [49] P. Moskal, Acta Phys. Polon. B **47**, 97 (2016)
- [50] P. Moskal, M. Skurzok and W. Krzemien, AIP Conf. Proc. **1753**, 030012 (2016)
- [51] M. Skurzok *et al.*, EPJ Web of Conferences **199**, 01018 (2019)
- [52] M. Skurzok *et al.*, arxiv:1908.03429 (2019)
- [53] O. Rundel, Doctoral Thesis, Jagiellonian University 2019, arxiv: 1905.04544s
- [54] M. Skurzok *et al.*, EPJ Web of Conferences **181**, 01014 (2018)
- [55] O. Rundel *et al.*, Acta Phys. Polon. B **48**, 1807 (2017)
- [56] O. Rundel *et al.*, EPJ Web Conf. **130**, 02008 (2016)
- [57] A. Khreptak, O. Rundel and M. Skurzok, EPJ Web of Conferences **199**, 05026 (2019)