

# Oxygen sensing ability of positronium

Saturday, 9 October 2021 12:10 (20 minutes)

Positronium (Ps) is an exotic atom consisting of a positron and an electron, and around  $10^{11}$  Ps atoms form in the human body during a PET scan. We have discussed little Ps in PET because its formation does not change the spatial information obtained by PET; Most Ps annihilates into back-to-back gamma-ray photons. However, Ps can provide other unique information due to the delay of the gamma-ray emission as long as its lifetime. The lifetime varies according to the chemical and physical environment for Ps. For example, the lower dissolved oxygen concentration ( $pO_2$ ), the longer a Ps atom survives in solutions. This is because of the unpaired electrons in the  $O_2$  molecule that enhance Ps annihilation via the electron exchange interaction. Knowing  $pO_2$  distribution is important for cancer patients because hypoxic cells are often resistant to radiotherapy as well as chemotherapy.

Moskal *et al.* named the new concept of PET as “Ps imaging [1],” and they also found that the Ps lifetime differs between healthy and cancer cells [2]. The difference may come from a combination of several chemical and physical conditions, but how each factor changes the Ps lifetime is a challenging matter to be understood. Some efforts have revealed how  $O_2$  molecules reduce Ps lifetime. Stepanov [3] found a positive correlation between the  $pO_2$  and Ps annihilation rate (the inverse of the lifetime) in water. Furthermore, we found good linearity between them by accumulating 50 times larger number of counts, as shown in Fig. 1 [4]. This line is namely calibration line for Ps as an oxygen sensor, and this result indicates a possibility of Ps as a hypoxia biomarker candidate. In other words, during a PET scan,  $10^{11}$  nanosized sensors for  $O_2$  are spontaneously created in vivo, and it is worth trying to read the indicator for improving cancer treatments.

**Fig. 1**  $pO_2$  vs Ps decay rate: squares (Lee [5]), triangles (Stepanov [3]), and circles (Shibuya [4]). (see at <https://www.nature.com/articles/s42005-020-00440-z/figures/3>)

## References

- [1] P. Moskal, *et al.*, *Nature Rev. Phys.*, **1**, 527–9 (2019).
- [2] P. Moskal, *et al.*, *bioRxiv* (2021). (doi: 10.1101/2021.08.05.455285)
- [3] P. S. Stepanov, *et al.*, *Phys. Chem. Chem. Phys.* **22**, 5123 (2020).
- [4] K. Shibuya, *et al.*, *Commun. Phys.*, **3**, 173 (2020).
- [5] J. Lee, *et al.*, *J. Chem. Phys.* **44**, 2506 (1966).

**Primary author:** SHIBUYA, Kengo (University of Tokyo, Japan)

**Presenter:** SHIBUYA, Kengo (University of Tokyo, Japan)

**Session Classification:** Saturday Noon Session