4th Jagiellonian Symposium on Advances in Particle Physics and Medicine



Contribution ID: 379

Type: poster

CP Discrete Symmetry study in the decay of ortho-Positronium atom using the J-PET detector.

Tuesday, 12 July 2022 11:01 (2 minutes)

Kavya Valsan Eliyan, Juhi Raj, On Behalf of the J-PET Collaboration Jagiellonian University.

Faculty of Physics, Astronomy and Applied Computer Science, Jagiellonian University, Kraków, Poland. Theranostics Center, Jagiellonian University, Kraków, Poland. Email: kavya.eliyan@doctoral.uj.edu.pl, juhi.raj@doctoral.uj.edu.pl

Interaction between electron-positron pair leads to direct annihilation into photons or creation of a bound state called Positronium. Positronium is the lightest purely leptonic object decaying into photons [1-3]. Positronium atom can be formed in two states based on the spin alignment of its constituting particles, Singlet state (1S0 -para-Positronium (p-Ps) and Triplet state (3S1- ortho-Positronium (o-Ps). Constrained by conservation laws, the o-Ps annihilate into odd number of photons (o-Ps \rightarrow 3 γ , where $\gamma = 1, 2, ...$), while the p-Ps decay into an even number of photons (p-Ps \rightarrow 2 γ , where $\gamma - 1,2, 3, ...$) [4,5]. As an atom bound by a central potential, it is a parity eigenstate, and as an atom built out of an electron and an anti-electron, it is an eigenstate of the charge conjugation operator [1]. Therefore, the positronium form o-Ps annihilation [6]. The Standard Model predicts that the photon–photon interaction and weak interactions will mimic the symmetry violation in the order of 10–9 and 10–13 respectively [6]. Violation of CP invariance in purely leptonic systems has never been seen so far [7]. The experimental limits on CP and CPT symmetry violation in the decays of o-Ps are set at the level of 10–3 [2,8].

In the year 2021, the limitations of the previous experiments were overcome by the J-PET detector due to its much higher granularity and improve the world result by a factor of three and reaches the statistical precision of 10-4. The reported result is the present best upper limit on the CP violation in the decay of ortho-Positronium, leaving us 5 orders of more statistical sensitivity to be explored in this aspect. J-PET detector is constructed of 192 polymer scintillators, where each scintillator is attached with photomultipliers at each end. 192 scintillators are arranged co-axially in three layers at 3 different radii 42.5 cm, 46.75 cm, 57.5 cm respectively. Positronium atom can be formed in the center of J-PET detector using the beta-emitter 22Na source placed inside a small chamber. The source is sandwiched between an aerogel material. Plastic scintillators offer high time and angular resolution. Time Over Threshold is adopted as a measure of energy deposition. The signals are measured by using the trigger-less data acquisition [9-12]. All of the previous investigations with Positronium, which tested the discrete symmetries, were based on symmetry odd operators constructed as the products of photons momenta ($k \boxtimes i$) and Positronium spin (S \boxtimes) vectors [2-4,6]. This project describes an extended study using another proposed operator [4], taking advantages of properties of the 3 layered J-PET detector, which enables to determine the linear polarization direction of annihilation photons. Measurement of polarization direction of annihilation photons (511 keV) is a unique feature of the J-PET detector which allows the study of CP symmetry violation by determining the expectation values of the CP symmetry odd operator ($\epsilon \boxtimes i \cdot k \boxtimes j$) where $i \neq j$). As a future prospect, the J-PET collaboration has developed a modular version of the J-PET detector to improve the detection efficiency of this measurement and provide larger statistics in a shorter duration of measurement time to improve the precision significantly.

ACKNOWLEDGEMENTS:

The author acknowledge support by the TEAM POIR.04.04.00-00-4204/17 program, the NCN grant no. 2021/42/A/ST2/00423 and the SciMat and qLife Priority Research Areas budget under the program Excellence Initiative - Research University at the Jagiellonian University.

References:

- [1] M. Skalsey, J. Van House, Phys. Rev. Lett. 67, 1993 (1991).
- [2] P. A. Vetter, S.J. Freedman, Phys. Rev. Lett. 91, 263401 (2003).
- [3] T. Yamazaki, T. Namba, S. Asai, T. Kobayashi, Phys. Rev. Lett. 104, 083401 (2010).
- [4] P. Moskal et al., Acta Phys. Polon. B 47, 509 (2016).
- [5] E.Czerwinski et al., Acta Phys. Polon. B 48, 509 (2017).
- [6] P. Moskal, A. Gajos, M. Mohammed, Nat. Commun. 12, 5658 (2021).
- [7] V. A. Kostelecky and N. Russell, 2018 update to Rev. Mod. Phys. 83, 11 (2011).
- [8] T. Yamazaki, T. Namba, S. Asai, T. Kobayashi, Phys. Rev. Lett. 104, 083401 (2010).
- [9] M. Palka, Jour. Of Instr. 12, P08001 (2017)
- [10] G. Korcyl et al., Acta Phys. Polon. B 47, 491 (2016).
- [11] Sz. Niedzwiecki et al., Acta Phys. Polon. B 48, 1567 (2017).
- [12] A. Gajos et al., Nucl. Instr. and Meth. A 819, 54 (2016).

Presenter: VALSAN ELIYAN; JAGIELLONIAN UNIVERSITY, POLAND, Kavya

Session Classification: Poster session & Coffee & Conference photo I